Self-Tanning Formulations are Improved with the Addition of Floraesters K-20W Jojoba

Presented by: Tiffany N. Oliphant, M.S., C.C.R.C. and Robert A. Harper, Ph.D.

For the Society of Cosmetic Chemists 2013 Sunscreen Symposium

September 19th - 21st, 2013 Orlando, FL



Contact: sales@floratech.com

Self-Tanning Formulations are Improved with the Addition of Floraesters® K-20W Jojoba

Tiffany N. Oliphant, M.S., C.C.R.C. (Floratech, Chandler, AZ) and Robert A. Harper Ph.D. (Harper & Associates, La Jolla, CA) Email: sales@floratech.com Website: www.floratech.com

Abstract

Data from multiple, pilot, clinical studies have shown that Floraesters K-20W Jojoba (K-20W) enhances the efficacy and sensory properties of multiple finished cosmetic and personal care formulations. For example, K-20W in combination with glycerin enhances skin moisturization; sunscreen actives in combination with K-20W are retained on the skin after water immersion; and some fragrances are retained on the skin longer in the presence of K-20W. Theoretically, the activity is due to the film-forming ability of the K-20W. In addition to these studies we now show that K-20W can also prolong the artificial skin color produced by a sunless tanner (i.e. self-tanner), while also improving its sensory appeal to consumers.

In a small, double-blind, vehicle-controlled, clinical study, subjects' backs were treated with one application of several sunless tanner formulations, all containing 5% DHA.1 Formulations varied with regard to pH and the amount of K-20W, acrylates/octylacrylamide copolymer,² and erythrulose.³ Skin color and moisturization measurements were taken before treatment, and 24, 48, 72, and 96 hours after treatment. Test articles containing K-20W (at 1%), or a combination of K-20W and erythrulose (at 0.5-1%), produced the greatest percent increase and retention of skin color (p<0.05 compared to the vehicle) at all time points. Test articles containing K-20W also produced higher levels of skin moisturization 24 hours after test article application. Test articles with and without 0.5% K-20W were also evaluated in consumer preference studies on the legs with repeat applications. Consumers preferred the formulations that contained K-20W for evenness of tan, longevity of tan, moisturization, odor, and overall preference. These results show that the addition of Floraesters K-20W Jojoba into sunless tanner formulations can enhance the effectiveness of these safe alternatives to sun exposure.

Background Information

- Floraesters K-20W Jojoba [INCI: Hydrolyzed Jojoba Esters (and) Water (Aqua)] is composed of 20% hydrolyzed jojoba esters and 80% water. Hydrolyzed jojoba esters are derived from jojoba (Simmondsia chinensis) seed oil.
- Dihydroxyacetone is utilized within sunless tanner formulations at 3-5%. DHA is a sugar that interacts with amino acids in the skin to result in a brown color change (stain) which lasts several days.4 According to the Academy of Dermatology, sunless tanning formulations containing DHA are the most effective.5
- Erythrulose is also a sugar used in sunless tanning formulations. It utilizes the same mechanism as DHA, however the reaction occurs more slowly thereby extending the life of the tan.6
- Dermacryl-79 (INCI: acrylates/octylacrylamide copolymer) is a film-forming polymer used to maintain active ingredients at the site of application.7

References / Footnotes

- Dihydroxyacetone (DHA) was supplied by EMD Chemicals Inc.
- Dermacryl-79 (INCI: Acrylates/Octylacrylamide Copolymer) was supplied by Akzo Nobel Chemicals
- Erythrulose was supplied by DSM Nutritional.
- SB Levy, Dihydroxyacetone-containing sunless or self-tanning lotions. Journal of the American Academy of Dermatology. 27(6): 989-993. 1992.
- http://www.medicinenet.com/script/main/art.asp?articlekey=23898 Accessed August 6, 2013. Web.
- CG Burkhart and CN Burkhart. Dihydroxyacetone and methods to improve its performance as artificial tanner. The Open Dermatology Journal. 3: 42-43. 2009. http://sc.akzonobel.com/en/personalcare/Pages/product-detail.
- aspx?prodID=6638 Accessed August 6, 2013. Web. Mexameter MX 18 is a product of Courage+Khazaka Electronic GmbH
- Corneometer is a registered trademark of Courage + Khazaka Electronic
- (Koln, Germany). GmbH (Koln, Germany).

Increased Color Retention

Objective: To determine if Floraesters K-20W Jojoba could increase skin color retention when incorporated into a sunless tanning formulation which contained 5% DHA. Design: One application of each of the sunless tanning formulations was made (2.5 mg/ cm²) to randomized locations on the lower backs of fifteen male and female subjects. End Point: Melanin measurements (i.e. skin color) via Mexameter MX 188 were conducted at baseline, and 24, 48, 72, and 96 hours, post sunless tanning formulation application.

Figure 1: Color Retention




Figure 1. Floraesters K-20W Jojoba enhanced skin color retention in a sunless tanning formulation better than (p<0.05) the vehicle test articles at all measured time points.

Increased Skin Hydration

Objective: To determine if Floraesters K-20W Jojoba (in conjunction with glycerin) could increase skin hydration when incorporated into a sunless tanning formulation which contained 5% DHA.

Design: One application of each of the sunless tanning formulations was made (2.5 mg/ cm²) to randomized locations on the lower backs of fifteen male and female subjects. End Point: Skin hydration measurements via Corneometer® CM 8259 were conducted at baseline and 24 hours post sunless tanning formulation application.




Figure 2. Floraesters K-20W Joioba increased skin hydration in a sunless tanning formulation containing 5% dihydroxyacetone better than (p<0.05) all other test articles at 24 hours.

Self-Tanning Formulations are Improved with the Addition of Floraesters® K-20W Jojoba

Tiffany N. Oliphant, M.S., C.C.R.C. (Floratech, Chandler, AZ) and Robert A. Harper Ph.D. (Harper & Associates, La Jolla, CA) Email: sales@floratech.com Website: www.floratech.com


Enhanced Consumer Preference

Objective: To determine the consumer preference between a sunless tanning formulation with and without Floraesters K-20W Jojoba.

Design: Randomized daily applications (2.5 mg/cm²) of two sunless tanner formulations, for three days, to the outer lower legs of 27 female subjects.

End Point: A consumer preference survey was completed on Day 3 for immediately observable skin and product properties and on Day 7 for long-term properties.

Figure 3: Product Characteristics




Evaluation	Day
Smell of Product	3
Spreadability	3
Slide Across Skin	3
Rinse-Off (hands)	3
Moisturization	3
Overall Product Preference	7
Overall Color	7
Evenness of Tan	7
Longevity of Tan	7
Overall Tanning Experience	7

Figure 4: Skin Characteristics

Overall Tanning Experience



Figures 3 and 4. The sunless tanner formulation with Floraesters K-20W Jojoba was preferred by consumers over the vehicle test article.



Conclusions

- Floraesters K-20W Jojoba increased skin color retention when incorporated into a sunless tanning formulation.
- Floraesters K-20W Jojoba increased skin hydration when incorporated into a sunless tanning formulation.
- 80% of consumers preferred the smell of the sunless tanning formulation with Floraesters K-20W Jojoba.
- Floraesters K-20W Jojoba increased consumer perception when incorporated into a sunless tanning formulation.

iLabel® - Floratech Information Instantly

iLabel delivers valuable product information instantly. It supplies lot-specific information including certificates of analysis, material safety data sheets, product specifications and technical data.

iLabel saves time by providing instant access to global regulatory information, clinical efficacy data, and demonstration formulas featuring Floratech products.

iLabel is an easy-to-use tool that does not require registration, usernames, or passwords.

